

Review of Recent Jlab Results

Elton S. Smith

Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

Abstract. High quality polarized electron beams at Jefferson Lab make possible precision measurements of hadronic properties in the regime of strongly interacting QCD. We will describe a few programs at Jefferson Lab that are making measurements that link the basic static properties of hadrons to their quark sub-structure. For example, parity-violating electron proton elastic scattering probes the spatial distribution of strange quarks in the nucleon. The nucleon-Delta transition form factors give us information about the deformation of nucleons and Deltas. Finally, new high statistics measurements of photons scattering off proton and deuteron targets are used to set upper limits on the production of exotic baryons with strangeness $S=+1$. These examples will be used to illustrate the capabilities and focus of the experimental program at JLab.

INTRODUCTION TO JLAB

The Continuous Electron Beam Accelerator Facility (CEBAF) at the Thomas Jefferson National Accelerator Facility (Jefferson Lab) is devoted to the investigation of the electromagnetic structure of mesons, nucleons, and nuclei using high energy and high duty-cycle electron and photon beams.

CEBAF is a superconducting electron accelerator [1] with a maximum energy of 6 GeV and 100 % duty-cycle. Three electron beams with a maximum total current of 200 μA can be used simultaneously for electron scattering experiments in the experimental areas. The accelerator design concept is based on two parallel superconducting continuous-wave linear accelerators joined by magnetic recirculation arcs. The accelerating structures are five-cell superconducting niobium cavities with a nominal average energy gain of 5 MeV/m. The accelerator operation has met all design goals, achieving 5.7 GeV for physics running, and delivering three high quality beams with intensity ratios exceeding $10^6:1$. The electron beam is produced using a strained GaAs photocathode allowing the delivery of polarized electrons ($P_e \geq 75\%$) simultaneously to all three halls.

Three experimental areas are available for simultaneous experiments, the only restriction being that the beam energies have to be multiples of the single pass energy. The halls contain complementary equipment which cover a wide range of physics topics: Hall A [2] has two high resolution magnetic spectrometers with 10^{-4} momentum resolution in a 10% momentum bite, and a solid angle of 8 msr. Hall B houses the large acceptance spectrometer, CLAS [3]. Hall C uses a combination of a high momentum spectrometer (10^{-3} momentum resolution, 7 msr solid angle and maximum momentum of 7 GeV/c) and a short orbit spectrometer.

To illustrate the physics which is being addressed at Jefferson Lab, we have chosen three topics of current interest: measurements of parity violation in electron elastic

scattering, the N- Δ transition form factors, and high statistic searches for the Θ^+ exotic strangeness +1 baryon with the CLAS. These programs have the common goal of probing baryon structure which comes from quark pairs beyond the contribution from the standard three valence quarks.

STRANGE QUARKS IN THE NUCLEON

Early deep-inelastic scattering experiments demonstrated that the structure of the proton can not be described by its uud valence structure alone. For example, quarks carry only 50% of the proton momentum and are responsible for less than 30% its spin [4]. The fraction and distribution of s and $\bar{s}$ quarks were also determined in ν and $\bar{\nu}$ scattering experiments. For example, NuTeV [5] reports the fraction $(s + \bar{s})/(\bar{u} + \bar{d})$ to be $0.42 \pm 0.07 \pm 0.06$ at a Q^2 of 16 GeV². But most of these sea quarks are very short lived ($\sim \hbar/\sqrt{Q^2}$) and arise from perturbative evolution of QCD [6]. The focus of this section is to describe experiments that are sensitive to the long-lived component of $s\bar{s}$ pairs which contribute to the static properties of the nucleon such as its magnetic moment and charge radius.

It is easy to understand how $s\bar{s}$ pairs in the sea can contribute to the proton magnetic moment in a simple hadronic picture [7]. During the process of $uuds\bar{s}$ fluctuations, the quarks will tend to arrange themselves into energetically favorable configurations, the lowest state being a ΛK^+ . The K^+ must be emitted in an L=1 state to conserve angular momentum and parity. The probability that the Λ has its spin anti-parallel to the proton is twice as likely as the probability that the spin have the same direction. In this case, the (positive) kaon will have $l_z = +1$ and contribute a positive amount to the magnetic moment of the proton. In addition, the spin of the (negative) s quark in the Λ is anti-aligned with the proton spin so it also gives a positive contribution to the magnetic moment. By convention, however, the usual definition of the strange magnetic moment ($G_M^s(Q^2 = 0)$) does not include the s -quark charge of $(-\frac{1}{3})$, and therefore this picture actually predicts a *negative* strange magnetic moment. Over the past decade there have been many different models [8] used to estimate both the magnitude and sign of G_M^s , but all have been forced to make non-trivial approximations. Most models predict a negative G_M^s , as in the case of the naive hadronic model above, with a value in the range of -0.8 to 0 . But it is clear for the moment that experimental measurements are required to guide our understanding of the contributions of strange quarks to the properties of the proton.

Parity violation in electron scattering is a unique tool for probing the contribution of $s\bar{s}$ sea quarks to the structure of the nucleon. The weak interaction is probed by measuring the electron helicity dependence of the elastic scattering rate off unpolarized targets. Experimental sensitivities of the order of a part per million are required to measure the parity violating asymmetry on a nucleon which can be expressed as

$$A_{PV}^p = \frac{d\sigma_R - d\sigma_L}{d\sigma_R + d\sigma_L} = \frac{G_F Q^2}{4\pi\alpha\sqrt{2}} \left[\frac{A_E + A_M + A_A}{\epsilon (G_M^\gamma)^2 + \tau (G_M^\gamma)^2} \right], \quad (1)$$

where $A_E = \varepsilon G_E^Z(Q^2) G_E^\gamma(Q^2)$, $A_M = \tau G_M^Z(Q^2) G_M^\gamma(Q^2)$ are the electric and magnetic γ -Z interference terms, $\tau = Q^2/4M_N^2$, and ε is the polarization of the virtual photon. The isolation of the electric and magnetic terms can be accomplished by taking advantage of the sensitivity to kinematics (though ε and τ) and taking data at different scattering angles. The last term, A_A , picks out the contribution due to the axial form factor, but the forward-angle measurements are insensitive to this term, so we refer the reader to our references for details [9]. Finally, the flavor decomposition of the form factors can be determined uniquely assuming charge symmetry and by combining parity-violating asymmetries with measurements of the electric and magnetic form factors on the proton and neutron. The scattering amplitude from a zero spin isoscalar target, such as ${}^4\text{He}$, is particularly simple because it does not allow magnetic transitions so the asymmetry is sensitive only to the isoscalar electric form factor:

$$A_{PV}^{He} = \frac{G_F Q^2}{4\pi\alpha\sqrt{2}} \left[4\sin^2\theta_W + \frac{G_E^s}{\frac{1}{2}(G_E^{\gamma p} + G_E^{\gamma n})} \right], \quad (2)$$

The contribution of the asymmetry which is attributable to strange quarks is obtained by subtracting the calculated asymmetry assuming no strange contribution, including radiative corrections and best estimates of the electromagnetic form factors. The uncertainties in the calculations are estimated and included separately when quoting the uncertainties in the strange form factors. Thus a fairly extensive experimental program is required to isolate all contributions.

There are two major experimental programs at JLab studying parity violation: HAPPEX [14] and G0 [15]. Both collaborations have recently reported results. The precision of these experiments, together with other experimental and theoretical efforts world-wide [9, 16, 17, 18, 19, 20], has now achieved sufficient accuracy to be sensitive to the contribution of strange quarks to the nucleon form factors. The first measurements of parity-violating electron scattering at JLab were carried out by the HAPPEX collaboration. The experiment detected the scattered electrons in each of the Hall A high resolution spectrometers, with sufficient hardware resolution to spatially separate the elastically scattered events from inelastic events into special lead-scintillator absorption counters whose output was integrated during a 30 ms window. Window pairs of opposite helicity were chosen randomly at 15 Hz. A half wave plate in the CEBAF injector was inserted or removed about once a day, switching the helicity of the beam without changing the electronics. HAPPEX has previously reported measurements of parity violation in ep elastic scattering at $Q^2=0.48 \text{ GeV}^2$ [21]. The asymmetry errors are completely dominated by statistical uncertainties, and the helicity correlated systematic uncertainties are about 0.1 ppm. Recently it has completed asymmetry measurements at $Q^2=0.1 \text{ GeV}^2$ in ep scattering giving $G_E^s + 0.080 G_M^s = 0.030 \pm 0.025 \pm 0.006 \pm 0.012$ [22] and a measurement of elastic scattering on ${}^4\text{He}$ at a $Q^2=0.091 \text{ GeV}^2$ which yields a direct measurement of $G_E^s = -0.038 \pm 0.042 \pm 0.010$, consistent with zero. [23]. These measurements were performed at electron scattering angles of 6° , made possible by the new superconducting septum magnets installed in Hall A [24]. We note that HAPPEX continues to take data and is expected to improve the statistical uncertainty in each measurement by a factor of 2.5–3 this year.




FIGURE 1. Constraints on the G_E^s and G_M^s strange form factors at $Q^2 = 0.1 \text{ GeV}^2$. The curves give the 95% contours. Sample model calculations are also indicated as 1 [10] 2 [11] 3 [12] and 4 [13].

The G0 experiment uses a dedicated detector [25] to determine G_E^s , G_M^s , and G_A^e over a broad kinematical range. To date measurements have been completed for forward angles. The result of their first experimental run has been the determination of the linear combination $G_E^s + \eta G_M^s$ for Q^2 between 0.12 to 1.0 GeV^2 ($\eta = \tau G_M^p / \epsilon G_E^p$). These measurements are consistent with HAPPEX and A4, but they provide broad coverage in Q^2 to probe the spatial dependence of strange quarks in the nucleon. The three lowest Q^2 points (out of 16 measurements) can be used together with the other experiments (see Fig. 1) to determine a central value for G_E^s and G_M^s separately for $Q^2 = 0.1 \text{ GeV}^2$.

We close by summarizing the experimental situation, and relating the measured quantities back to our simple hadronic picture for possible strange contributions to the proton magnetic moment. Analysis of the world measurements leads to $G_M^s = 0.55 \pm 0.28$ and $G_E^s = -0.01 \pm 0.03$. To compare to the proton magnetic moment of $+2.79 \text{ n.m.}$ we multiply G_M^s by $(-\frac{1}{3})$ and, taken at face value, it is approximately 7% of the magnitude of the proton magnetic moment, and opposite in sign. The interpretation of positive values for G_M^s has been investigated within the context of the quark model [26]. This study shows that, even though the quark model is very successful at predicting relations between baryon magnetic moments, the quark model does not lead naturally to positive values for G_M^s . The authors find that positive moments are generated when the $\bar{s}$ is in the ground state and the $uuds$ quarks are in excited states. Although these configurations do not seem very natural, they are very analogous to quark-models configurations for pentaquarks, which is the topic of the last section of this paper.

N- Δ DEFORMATION

Within the spin-flavor SU(6) symmetry of the quark model, the Δ resonance is a completely symmetric object with all quark spins aligned. Within this model, the photo-excitation $\gamma^* N \rightarrow \Delta(1232) \rightarrow N\pi$ proceeds via a single quark spin flip in the L=0 nucleon ground state. This magnetic dipole transition, characterized by the M_{1+} multipole, predicts negligible contributions from the electric (E_{1+}) and scalar (S_{1+}) quadrupole multipoles. However, strong-interaction dynamics will modify this picture, and several models predict deviations from this simple picture which would indicate deformations of both the nucleon and the Δ [27]. For example, the deformation of the nucleon and Δ could result from D-state admixtures into the baryon wave function. However, D-wave admixtures are expected to be small, but quark-antiquark pairs which are present in the nucleon can also contribute to the quadrupole moment. These latter contribute to the quadrupole moment via a two-body spin tensor in the charge operator, even when the valence quarks are in pure S states. The physical interpretation in this case is that the distribution of $q\bar{q}$ sea quarks in the nucleon deviate from spherical symmetry. These models interpret the negative values of the measured transition quadrupole moments as corresponding to a positive intrinsic quadrupole moment (cigar shape) for the nucleon and a negative moment (pancake shape) for the Δ .

Complete angular distributions of the pion in the reaction $ep \rightarrow ep\pi^0$ at the peak of the Δ allow the experimental determination of the quadrupole moments. The differential cross section of a polarized beam (helicity $h = \pm 1$) and unpolarized target can be written as a function of the transverse and longitudinal structure functions σ_T and σ_L , and the interference terms σ_{TT} , σ_{LT} and $\sigma_{LT'}$:

$$\begin{aligned} \frac{d^2\sigma}{d\Omega_\pi^*} = & \frac{p_\pi^*}{k_\gamma^*} \left(\sigma_T + \varepsilon_L \sigma_L + \varepsilon \sigma_{TT} \sin^2 \theta_\pi^* \cos 2\phi_\pi^* \right. \\ & \left. + \sqrt{2\varepsilon_L(\varepsilon+1)} \sigma_{LT} \sin \theta_\pi^* \cos \phi_\pi^* + h \sqrt{2\varepsilon_L(\varepsilon-1)} \sigma_{LT'} \sin \theta_\pi^* \sin \phi_\pi^* \right), \end{aligned} \quad (3)$$

where p_π^* , θ_π^* , and ϕ_π^* are the pion center-of-mass (c.m.) momentum and angles, ε and ε_L are the usual transverse and longitudinal polarizations of the virtual photon, and k_γ^* is the real-photon-equivalent c.m. energy. The moments of interest can be selected as terms in the partial wave expansion of the structure functions using Legendre polynomials [28, 29]. Unique solutions can be obtained by truncating the multipole expansion to terms involving only M_{1+} . Fitting for the coefficients of the expansion allows the extraction of the electric (E_{1+}) and scalar (S_{1+}) quadrupole moments through their interference with the dominant M_{1+} . The systematic errors due to the truncation of higher multipoles are estimated by calculating the effects of higher partial waves using realistic parameterizations of higher mass resonances and backgrounds. At the peak of the Δ and for Q^2 of a few GeV^2 , this procedure results in a largely model-independent extraction of the moments.

Early interest in the quadrupole moments stemmed from the perturbative QCD predictions of quark helicity conservation which requires that $E_{1+} \rightarrow M_{1+}$ and S_{1+} approaches a constant at very high Q^2 . Therefore, the first experiments at JLab in Hall C [30] determined the quadrupole moments at the highest accessible $Q^2 \sim 3 - 4 \text{ GeV}^2$, followed




FIGURE 2. Q^2 dependence of the ratios of the electric (E_{1+}) and scalar (S_{1+}) quadrupoles to the dominant dipole moment. The curves are model calculations from references [32], [33] and [34].

by results from CLAS [31] at intermediate momentum transfer. These measurements showed that E_{1+} was not only a few percent of M_{1+} , but also of opposite sign. The asymptotic kinematics of perturbative QCD was clearly well beyond the range of these measurements.

Recently the focus has shifted to understanding the deformation of the nucleon at very low Q^2 which probes chiral symmetry breaking, models of heavy baryon chiral perturbation theory, and the shape of the pion cloud surrounding the nucleon. Data at $Q^2 = 0.12 \text{ GeV}^2$ from MAMI [35] and Bates [36] show a surprisingly large scalar quadrupole moment of 6.5%, twice that expected from the trend of higher energy measurements and most models. New data from CLAS [37] were taken with an electron beam energy of 1.046 GeV, polarization of $\sim 70\%$ and 10 nA beam current on a 2-cm liquid hydrogen target. The π^0 electroproduction data from this experiment have provided nearly complete angular distributions for $Q^2 = 0.14 - 0.38 \text{ GeV}^2$ and $W = 1.2 - 1.3 \text{ GeV}$. The broad acceptance of the CLAS detector allows measurements of the full angular distribution over a broad range of Q^2 and covering the W range of the Δ . The ratios of E_{1+}/M_{1+} and S_{1+}/M_{1+} from this experiment along with other measurements are shown in Fig. 2. Important constraints on the non-resonant backgrounds have been obtained by measuring the helicity-dependent structure function $\sigma_{LT'}$ [38]. Weak non-resonant backgrounds underlying the Δ peak can be enhanced through their interference with the magnetic multipole ImM_{1+} , which is suppressed in σ_{LT} due to the vanishing of

the real part of the resonant amplitude at the pole.

In summary, there is general agreement on the value of $E_{1+}/M_{1+} \sim -2.5\%$, which is approximately constant at small Q^2 . The recent analysis of CLAS indicates that S_{1+}/M_{1+} ratio shows a Q^2 -dependence in this same kinematic region, in apparent disagreement with previous measurements that show it is leveling off. Global parameterizations of electroproduction data is underway and will be used along with π^+ data to extract the nucleon parameters with refined accuracy.

SEARCH FOR PENTAQUARKS

Pentaquark states have been studied both theoretically and experimentally for many years [39]. Recent interest was revived by predictions within the chiral soliton model [40] for the existence of an anti-decuplet of 5-quark resonances with spin and parity $J^\pi = \frac{1}{2}^+$. The lowest mass member of the anti-decuplet, now called the Θ^+ , is an isosinglet with valence quark configuration $uudd\bar{s}$ with strangeness $S = +1$, was predicted to have a mass of $1.53 \text{ GeV}/c^2$ and a width of $\sim 0.015 \text{ GeV}/c^2$. These definite predictions prompted experimental searches to focus attention in this mass region. Shortly after the first observation of an exotic $S = +1$ state by the LEPS collaboration [41], many experiments presented confirming evidence of the initial report, including two observations using the CLAS detector [42, 43]. However, there have been an increasing number of experiments reporting null results, even though in very different reactions and disparate kinematic regions [44].

This paper reports on two searches for a narrow $S = +1$ baryon in the mass range between 1520 and 1600 MeV using the CLAS detector. These new high-statistics searches were conducted using the tagged photon beam facility in Hall B [45] with deuterium [46] and proton [47] targets. The deuterium data can be compared directly with the first reported observation by CLAS [42], which analyzed a data set referred to as “G2a.” The G2a data set consisted of two run periods at $E_e = 2.478 \text{ GeV}$ and 3.115 GeV , corresponding to 0.83 and 0.35 pb^{-1} respectively. The photon energy was tagged beginning below the reaction threshold of 1.51 GeV up to 95% of the electron beam energy. The target was 10 cm long and located on CLAS center, and the CLAS torus field was set at 90% of its maximum value.

The new deuterium data set, referred to as G10, used a primary electron beam energy of $E_e = 3.767 \text{ GeV}$ creating a tagged photon beam with energies E_γ between 1 and 3.6 GeV. The run was divided evenly between two field settings of the torus magnet (60% and 90% of the full-field current of 3860 A). The hardware trigger required two charged particles in any two sectors and the target was a 24-cm long cell containing liquid deuterium and located 25 cm upstream of nominal CLAS center. The integrated luminosity for this data set was 38 pb^{-1} for E_γ greater than 1.5 GeV. Evidence for the exotic baryon was searched for using the reaction $\gamma d \rightarrow K^- K^+ pn$. The momentum of the charged particles was determined using magnetic analysis and their mass determined using time-of-flight techniques. The analysis selected a detected proton, K^+ and K^- in the final state, all originating from the same beam bucket. The neutron momentum and energy was reconstructed from the measured charge particle tracks and known energy




FIGURE 3. (Left) $MM(pK^-)$ distribution of the G10 data (red histogram) compared to the G2a distribution (black points with error bars). Selection cuts have been made on the G10 data set to best represent the experimental conditions of G2a and scaled for direct comparison. (Right) Distribution of the nK^+ mass spectrum for the G11 data. No evidence for narrow structures is apparent. The inset shows the mass distribution with selection cuts to reproduce the SAPHIR analysis [48]. Note: All results from CLAS are preliminary.

of the incident photon. The exotic Θ^+ baryon was searched for in the decay $\Theta^+ \rightarrow K^+ n$, where the K^+ uniquely identified the positive strangeness of the baryonic state. The mass of the $K^+ n$ system is shown in Fig. 3(left) for both the G2 and G10 data samples. The spectrum is relatively smooth and does not exhibit the peak found in the original smaller data set. The figure overlays the two spectra with selection cuts on G10 to mimic the G2a data set as closely as possible including the photon energy spectrum. If we use the G10 data as a background shape, the peak at 1.54 GeV in the G2 data set is consistent with a 3σ fluctuation. The new data show no indication of a peak in the $\Theta^+ \rightarrow K^+ n$ exotic $S=+1$ channel.

The measurement on the proton (G11) was conducted at lower photon energy and using a different reaction from the published CLAS result [43]. This new experiment used a primary electron beam energy $E_e = 4.0$ GeV to produce a tagged photon beam in the range of E_γ between 1.6 and 3.8 GeV. The target consisted of a 40-cm long cylindrical cell with liquid hydrogen. The data were taken over a period of 50 days, corresponding to an integrated luminosity of 70 pb^{-1} , which is an order-of-magnitude greater than previous experiments with tagged photons. Evidence for the $S = +1$ exotic baryon Θ^+ was searched for in the reaction $\gamma p \rightarrow \bar{K}^0 K^+ n$. The search was conducted for the baryon produced in association with the $\bar{K}^0$ and subsequent decay $\theta^+ \rightarrow K^+ n$. The positive strangeness of the baryon was uniquely tagged using the K^+ of the decay. The charged kaon, as well as the pions from the $\bar{K}^0 \rightarrow \pi^+ \pi^-$ decay, were reconstructed in the CLAS detector. The neutron momentum was reconstructed using energy and momentum

conservation and the known energy of the incident photon. The missing mass peak of the neutron was reconstructed to be 939 MeV with a width of 10 MeV. The $\Sigma^+ \rightarrow \pi^+ n$ and $\Sigma^- \rightarrow \pi^- n$ reactions are also present in this data sample and are reconstructed with an accuracy of less than 1 MeV and resolution of 3 MeV. These baryons, as well as the $\Lambda^*(1520)$ were excluded from the present analysis. After all cuts, the reconstructed nK^+ mass spectrum is shown in Fig. 3(right). The spectrum is smooth and structureless, with no indication of a peak near 1540 MeV where the Θ^+ was previously reported [48, 49]. An upper limit to the Θ^+ cross section of less than 1 nb was set based on assuming a t-exchange production mechanism for the reaction.

We conclude with a brief summary of the status of pentaquark measurements with CLAS. There are two published observations [42, 43] of the Θ^+ baryon in photon beam reactions on deuterium and proton targets. The measurement on deuterium has been repeated with higher statistics and the new data do not confirm the original observation. A new experiment to verify the production off the proton in the reaction $\gamma p \rightarrow \pi^+ K^- K^+ n$ for photon energies between 3 and 5.5 GeV has been approved [50], but has not yet received beam time. However, a recent high statistics experiment on the proton at lower energy [51] has searched for the narrow exotic baryon in the reaction $\gamma p \rightarrow \bar{K}^0 K^+ n$ and found no evidence for the state in the nK^+ mass range between 1520 and 1600 MeV.

ACKNOWLEDGMENTS

I would like to thank Alberto dos Reis and all the organizers of the conference for their gracious hospitality. I would also like to thank K. Paschke, D. Armstrong and D. Beck of the HAPPEX and G0 collaborations for making available the results on parity violation. From the CLAS collaboration I would like to express special thanks to K. Joo and L. Smith for help in preparation of materials on Delta production, and S. Stepanyan, R. De Vita and M. Battaglieri for discussions of the pentaquark search. The Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility for the U.S. Department of Energy under contract DE-AC05-84ER40150.

REFERENCES

1. C. Leemann, D. Douglas, and G. Krafft, *Annu. Rev. Nucl. Part. Sci.* **51**, 413 (2001).
2. J. Alcorn, et al., *Nucl. Instr. Meth. A* **522**, 294–346 (2004).
3. B. Mecking, et al., *Nucl. Instr. Meth. A* **503**, 513–553 (2003).
4. E. Hughes, and R. Voss, *Annu. Rev. Nucl. Part. Sci.* **49**, 303 (1999).
5. T. Adams et al. (NuTeV) (1999), [arXiv:hep-ex/9906038].
6. M. Burkardt, and B. Warr, *Phys. Rev. D* **45**, 958 (1992).
7. L. Hannelius, D. Riska, and L. Glozman, *Nucl. Phys. A* **665**, 353 (2000).
8. D. Beck, and B. Holstein, *Int. J. Mod. Phys. E* **10**, 1 (2001), [arXiv:hep-ph/0102053], and references therein.
9. E. Beise, M. Pitt, and D. Spayde, *Prog. Part. Nucl. Phys.* **54**, 289 (2005), [arXiv:nucl-ex/0412054], and references therein.
10. D. Leinweber, et al., *Phys. Rev. Lett.* **94**, 212001 (2005), [arXiv:hep-lat/0406002].
11. V. Lyubovitskij, et al., *Phys. Rev. C* **66**, 055204 (2002), [hep-ph/0207225].

12. R. Lewis, W. Wilcox, and R. M. Woloshyn, *Phys. Rev. D* **67**, 013003 (2003).
13. A. Silva, H.-C. Kim, and K. Goeke, *Phys. Rev. D* **65**, 014016 (2002).
14. K. Aniol et al. (HAPPEX Collaboration), *Phys. Rev. C* **69**, 065501 (2004).
15. D. Armstrong et al. (G0 Collaboration), *Phys. Rev. Lett.* **95**, 092001 (2005), [arXiv:nucl-ex/0506021].
16. F. Maas et al. (A4 Collaboration), *Phys. Rev. Lett.* **93**, 022002 (2004), [arXiv:nucl-ex/0401019].
17. F. Maas et al. (A4 Collaboration), *Phys. Rev. Lett.* **94**, 152001 (2005), [arXiv:nucl-ex/0412030].
18. D. Beck, and R. McKeown, *Annu. Rev. Nucl. Part. Sci.* **51**, 189 (2001).
19. K. Kumar, and P. Souder, *Prog. Part. Nucl. Phys.* **45**, S333 (2000).
20. The 2005 Joint Jefferson Lab/Institute for Nuclear Theory Workshop, Precision ElectroWeak Interactions Workshop (2005), http://www.wm.edu/physics/PreciseEW_2005.html.
21. K. Aniol et al. (HAPPEX Collaboration), *Phys. Lett. B* **509**, 211 (2001).
22. K. Aniol et al. (HAPPEX Collaboration) (2005), [arXiv:nucl-ex/0506011].
23. K. Aniol et al. (HAPPEX Collaboration) (2005), [arXiv:nucl-ex/0506010].
24. P. Brindza, et al., *IEEE Trans. on Applied Superconductivity* **11**, 1294 (2001).
25. P. Roos, *Eur. Phys. J. A* **24**, S2.59 (2005).
26. B. Zou, and D. Riska, *Phys. Rev. Lett.* **95**, 072001–1 (2005), [arXiv:hep-ph/0502225].
27. A. Buchmann, and E. Henley, *Phys. Rev. C* **63**, 015202 (2001), [hep-ph/0101027].
28. V. Burkert, and T. H. Lee, *Int. J. Mod. Phys. E* **13**, 1035 (2004), [nucl-ex/0407020], and references therein.
29. V. Burkert, *Prog. Part. Nucl. Phys.* **55**, 108 (2005), and references therein.
30. V. Frolov, et al., *Phys. Rev. Lett.* **82**, 45 (1999), [hep-ex/9808024].
31. K. Joo et al. (CLAS Collaboration), *Phys. Rev. Lett.* **88**, 122001 (2002), [hep-ex/0110007].
32. T. Sato, and T. Lee, *Phys. Rev. C* **63**, 055201 (2001), [nucl-th/0010025].
33. S. Kamalov, et al., *Phys. Rev. C* **64**, 032201(R) (2001), [nucl-th/0006068].
34. Z. Li, Y. Dong, and W. Ma, *J. Phys. G* **23**, 151 (1997).
35. T. Pospisil, et al., *Phys. Rev. Lett.* **86**, 2959 (2001), [nucl-ex/0010020].
36. C. Mertz, et al., *Phys. Rev. Lett.* **86**, 2963 (2001), [nucl-ex/9902012].
37. L. Smith, $\gamma^* p \rightarrow \Delta(1232) \rightarrow \pi N$ Results from CLAS (2005), contribution to the Japan-US Workshop on *Electromagnetic Meson Production and Chiral Dynamics*, Osaka, Japan, <http://www-nuclth.phys.sci.osaka-u.ac.jp/jp-usw/prog.html>.
38. K. Joo et al. (CLAS Collaboration), *Phys. Rev. C* **68**, 032201 (2003), [nucl-ex/0301012].
39. P. D. Group, *Phys. Lett. B* **170**, 289 (1986).
40. D. Diakonov, V. Petrov, and M. Polyakov, *Z. Phys. A* **359**, 305 (1997), [arXiv:hep-ph/9703373].
41. T. Nakano et al. (LEPS Collaboration), *Phys. Rev. Lett.* **91**, 012022 (2003), [arXiv:hep-ex/0301020].
42. S. Stepanyan et al. (CLAS Collaboration), *Phys. Rev. Lett.* **91**, 252001 (2003), [arXiv:hep-ex/0307018].
43. V. Kubarovsky et al. (CLAS Collaboration), *Phys. Rev. Lett.* **92**, 032001 (2004), [arXiv:hep-ex/0311046].
44. A. Dzierba, C. Meyer, and A. Szczepaniak, “Reviewing the Evidence for Pentaquarks,” in *J. Phys. Conf.*, **9**, 2005, p. 192, [arXiv:hep-ex/0412077], and references therein.
45. D. Sober, et al., *Nucl. Instr. Meth. A* **440**, 263–284 (2000).
46. K. Hicks, and S. Stepanyan, spokespersons for Investigation of Exotic Baryons States in Photoproduction Reactions with CLAS (2003), JLab Experiment E-03-113.
47. M. Battaglieri, R. De Vita, and V. Kubarovsky, spokespersons for Spectroscopy of Exotic Baryons with CLAS: Search for Ground and Excited States (2004), JLab Experiment E-04-021.
48. J. Barth et al. (SAPHIR Collaboration), *Phys. Lett. B* **572**, 127 (2003), [arXiv:hep-ex/0307083].
49. M. Ostrick, *Prog. Part. Nucl. Phys.* **55**, 337 (2005).
50. M. Battaglieri, et al., spokespersons for Study of Pentaquark States in Photoproduction off Protons (2004), JLab Experiment E-04-017.
51. M. Battaglieri et al. (CLAS Collaboration), Search for $\Theta^+(1540)$ pentaquark in high statistics measurement of $\gamma p \rightarrow \bar{K}^0 K^+ n$ at CLAS (2005), in preparation.